2 osteogenic differentiation of human adipose derived stem cells for osteochondral tissue 3 engineering applications 4 Dr. Liliana F. Mellor<sup>3,#</sup>, Dr. Rachel C. Nordberg<sup>1,#</sup>, Pedro Huebner<sup>2</sup>, Dr. Mahsa Mohiti-Asli<sup>3</sup>, 5 Michael A. Taylor<sup>3</sup>, Dr. William Efird<sup>4</sup>, Dr. Julia T. Oxford<sup>5</sup>, Dr. Jeffrey T. Spang<sup>4</sup>, \*Dr. Rohan A. 6 Shirwaiker<sup>2,3</sup> and \*Dr. Elizabeth G. Loboa<sup>1</sup> 7 # These authors contributed equally to this work 8 9 <sup>1</sup> College of Engineering, University of Missouri, Columbia, MO 65211 10 <sup>2</sup> Edward P. Fitts Department of Industrial and Systems Engineering, North Carolina State 11 University, Raleigh, NC 27695, USA 12 <sup>3</sup> Joint Department of Biomedical Engineering at University of North Carolina at Chapel Hill and 13 North Carolina State University, Raleigh, NC 27695, USA 14 <sup>4</sup> Department of Orthopaedics, University of North Carolina School of Medicine, Chapel Hill, NC 15 27599 16 <sup>5</sup> Biomolecular Research Center, Boise State University, Boise, ID 17 18 \*Co-corresponding authors: 19 Email address: egloboa@missouri.edu and rashirwaiker@ncsu.edu 20 Phone: 573-882-4378, 919-515-6416 21 Fax: 573-882-2490 22

Investigation of multi-phasic 3D-bioplotted scaffolds for site-specific chondrogenic and

1

#### **ABSTRACT**

1

2

3

4

5

6

7

8

9

10

11

12

13

14

15

16

17

18

19

20

21

Osteoarthritis (OA) is a degenerative joint disease that limits mobility of the affected joint due to the degradation of articular cartilage and subchondral bone. The limited regenerative capacity of cartilage presents significant challenges when attempting to repair or reverse the effects of cartilage degradation. Tissue engineered medical products are a promising alternative to treat osteochondral degeneration due to their potential to integrate into the patient's existing tissue. The goal of this study was to create a scaffold that would induce site-specific osteogenic and chondrogenic differentiation of human adipose derived stem cells (hASC) to generate a full osteochondral implant. Scaffolds were fabricated using 3D-bioplotting of biodegradable polycraprolactone (PCL) with either β-tricalcium phosphate (TCP) or decellularized bovine cartilage extracellular matrix (dECM) to drive site-specific hASC osteogenesis and chondrogenesis, respectively. PCL-dECM scaffolds demonstrated elevated matrix deposition and organization in scaffolds seeded with hASC as well as a reduction in collagen I gene expression. 3D-bioplotted PCL scaffolds with 20% TCP demonstrated elevated calcium deposition, endogenous alkaline phosphatase activity, and osteopontin gene expression. Osteochondral scaffolds comprised of hASC-seeded 3D-bioplotted PCL-TCP, electrospun PCL, and 3D-bioplotted PCL-dECM phases were evaluated and demonstrated site-specific osteochondral tissue characteristics. This technique holds great promise as cartilage morbidity is minimized since autologous cartilage harvest is not required, tissue rejection is minimized via use of an abundant and accessible source of autologous stem cells, and biofabrication techniques allow for a precise, customizable methodology to rapidly produce the scaffold.

- 23 **Keywords:** human adipose derived stem cells, osteochondral, osteogenic differentiation,
- 24 chondrogenic differentiation, 3D-printing
- 25 Running Head: 3D-bioplotted scaffolds for osteochondral tissue engineering

#### INTRODUCTION

Osteoarthritis (OA) is a degenerative joint disease that limits mobility of the affected joint due to the degradation of articular cartilage and changes in the adjacent subchondral bone. OA has been estimated to affect approximately 27 million people in the United States, and the prevalence is only expected to grow (1). There are significant challenges with treating cartilage and osteochondral degradation due to the limited regenerative capacity of articular cartilage, lack of understanding of the molecular mechanisms that trigger early changes in cartilage homeostasis, and lack of tools to detect the disease at early stages. To date, there is no cure for OA and therapeutic treatments are limited to controlling pain and inflammation of the joint.

Current clinical treatment options for cartilage repair include autologous chondrocyte implantation (ACI), microfracture, and mosaicplasty (2). ACI requires a surgical biopsy of the patient's healthy articular cartilage, followed by isolation and expansion of chondrocytes for several weeks in culture. A second surgery reintroduces the expanded chondrocytes into the defect site. However, this method has associated donor site morbidity and yields fibrocartilaginous tissue which is prone to degeneration (3). In microfracture, small holes are drilled into the defect to allow mesenchymal stem cells to migrate from the subchondral bone yielding new tissue formation (4). This technique also results in fibrocartilage formation (5), and therapeutic effects are only sustained for approximately one year (4). Mosaicplasty transfers plugs of autologous osteochondral tissue from non-weight-bearing healthy regions to repair the defect. However, this method is limited by the amount of donor sites and is also associated with donor site morbidity (6). Because of the limitations with current treatment options for osteochondral injury, tissue engineering approaches have become an active area of research.

One of the critical challenges from a clinical application perspective is creating an engineered tissue that can mimic the complex multi-phasic nature of osteochondral tissue (7). Because of

these challenges, many stem cell based approaches for cartilage repair have focused on cartilage regeneration (6, 8-16) instead of full osteochondral tissue regeneration (17-20). Biphasic osteochondral scaffolds have been generated by assembling two separate scaffolds of engineered bone and cartilage tissue (21-23). A disadvantage of this method is the lack of mechanical integration at the junction, leading to delamination of the layers. One approach to avoid delamination is to develop a monolithic scaffold with two phases that can be cultured as a unit to induce osteogenesis and chondrogenesis simultaneously. In this method, there are challenges to promoting both chondrogenic and osteogenic factors within the same culture system (24), which can be addressed by introducing biomimetic factors to the scaffold that induce either cartilage or bone formation (25, 26). In addition to biphasic systems, a few studies have attempted to achieve a graded transition zone of bone to cartilage throughout the depth of an engineered osteochondral tissue (26-31) as an attempt to improve load-bearing capabilities in vivo and reduce shear stresses at the biomaterial interfaces (32). Some of the technical and culturing challenges include identifying the ideal cell source, optimizing growth factors and signaling molecules to induce site-specific differentiation, inhibiting hypertrophy, and developing a simple and reliable manner to manufacture scaffolds of varying size to fit a specific injury site. After implantation, the primary concern is blood vessel invasion into the cartilage region that results in endochondral ossification, and proper integration with the host tissue (33).

19

20

21

22

23

24

25

26

1

2

3

4

5

6

7

8

9

10

11

12

13

14

15

16

17

18

The objective of this study was to investigate the use of a single cell source, human adipose derived stem cells (hASC) (34, 35), within an integrated multi-material, multi-scale triphasic 3D scaffold created using a combination of 3D-bioplotting and electrospinning to achieve complete osteochondral tissue generation. Human ASC, which are more abundant and accessible than human bone marrow derived mesenchymal stem cells (hMSC) (36-38), are capable of both osteogenic and chondrogenic differentiation and can express bone and cartilage extracellular matrix (ECM) constituents(38). Although some reports indicate that hASC chondrogenesis may

be inferior to hMSC chondrogenesis (39, 40), this can be overcome by alternative growth factor supplementation(41). Our triphasic 3D scaffold consists of physiologically inspired chemical cues to induce site-specific hASC differentiation in a manner that recapitulates the depth-dependent properties of native osteochondral tissue. The integrated 3D scaffold includes a 3D-bioplotted deep layer of a composite of polycaprolactone (PCL) and β-tricalcium phosphate (TCP) to induce osteogenesis, a 3D-bioplotted superficial layer of PCL combined with decellularized articular cartilage ECM (dECM) hydrogel to induce chondrogenesis, and an intermediate electrospun layer of PCL to mimic the native tidemark of osteochondral tissue to separate the two phases and potentially prevent blood vessel invasion into the cartilage layer *in vivo* (**Figure 1**).

#### MATERIALS AND METHODS

- 2 We first conducted analyses of hASC osteogenesis and chondrogenesis on separate phases of
- 3 3D-bioplotted PCL-TCP and PCL-dECM scaffolds, respectively, and then assessed the site-
- 4 specific hASC differentiation characteristics of integrated multi-material triphasic scaffolds
- 5 resembling full thickness osteochondral tissue.

6

7

1

# Preparation of PCL-TCP composite

- 8 The PCL-TCP composite (80-20% by weight) used for 3D-bioplotting of the osteogenic scaffold
- 9 phase was created by mixing the appropriate proportion of PCL (M<sub>n</sub>: 80,000, ~ 3 mm pellets;
- 10 Sigma-Aldrich, St. Louis, MO) and TCP (Riedel-de-Haën AG, Seelze, Germany), and heating
- the mixture at 180°C for 60 minutes with intermittent stirring in a temperature-controlled oven
- until a homogeneous, single phase mix was obtained.

13

14

# Preparation of decellularized ECM hydrogel

- 15 Decellularized ECM derived from bovine hooves (Micro Summit Processors, Micro, NC) was
- used in a hydrogel form as a constituent of the 3D-bioplotted scaffolds for the chondrogenic
- 17 phase. Cartilage was harvested from fresh hooves and decellularized by suspending in a
- 18 solution of deionized water and 0.5% penicillin/streptomycin and mixing for 24 hours. The
- 19 decellularized cartilage was then frozen in 1X PBS and lyophilized for 24 hours. Finally, the
- 20 lyophilized cartilage was pulverized in a mill using a number 40 size mesh to obtain the
- 21 decellularized cartilage ECM powder.

- ECM gels were prepared at an initial concentration of 25 mg/mL. The dECM was digested in a 2
- 24 mg/mL solution of pepsin in 0.1 M HCl. 4 mL of this solution was added to 100 mg of cartilage
- 25 dECM powder. The ECM was digested in the pepsin solution for 48 hours. Next, 400 μL of 1 M
- NaOH was added to the solution in order to bring the pH to 7.4. Finally, 489 µL of 10X PBS was

- added to the solution to make the concentrations of NaOH, HCI, and PBS equal to each other.
- 2 The final concentration of the gel was calculated to be 20.5 mg/mL.

# Proteomic analysis of decellularized ECM

Proteins were extracted using RIPA buffer protocol (Millipore, Billerica, MA). Protein concentrations were determined using a BCA protein assay (Thermo Fisher Scientific, Waltham, MA). 20 µg of total protein from each sample was digested in solution with Trypsin/Lys C mix (Promega) following the manufacturer's instructions. Briefly, protein samples were precipitated with cold acetone and dissolved in 8 M urea solution. After reduction with dithiothreitol and alkylation with iodoacetamide, samples were diluted with 50 mM Tris-HCl pH 8.0, to reduce urea concentration to 1 M. Trypsin/Lys C Mix was added at a 20:1 protein:protease ratio (w/w). Samples were digested overnight at 37 °C. Digestion was terminated with trifluoroacetic acid. Resulting peptide mixtures were cleaned using a C18 reverse-phase spin column (Thermo Fisher Scientific), dried under vacuum, and reconstituted in 5% acetonitrile, 0.1% formic acid in water for LC-MS/MS analysis.

LC-MS/MS analysis of peptides was conducted on a Velos Pro Dual-Pressure Linear Ion Trap mass spectrometer equipped with a nano electrospray ionization source and coupled with an Easy-nLC II nano LC system (Thermo Fisher Scientific). 5 μL of peptide mixture was loaded onto a C18 reverse-phase column (10 cm x 75 μm, 3 μm, 120 Å). Each sample was analyzed three times. A linear gradient with two mobile phases (A: 5% acetonitrile, 0.1% formic acid, 94.9% water, 0; B: 80% acetonitrile, 0.1% formic acid, 19.9% water) at a flow rate of 300 nL/min was used to separate peptide mixtures. The gradient began at 0% B, increased linearly to 50% B over 180 minutes and then to 100% B over 16 minutes, and maintained at this percentage for 14 minutes as a washing step. Eluted peptides were ionized in a nano ESI source with a spray voltage of 2.2 kV. Full scan MS spectra were acquired from m/z 300-2000. Collision-induced

dissociation (CID) was used to fragment the precursor ions. MS/MS spectra were acquired in

the data-dependent acquisition mode for the ten most abundant precursor ions in the preceding

full MS scan.

Peptide spectral matching and protein identification were achieved by database search using Sequest HT algorithms in Proteome Discoverer 1.4 (Thermo Fisher Scientific). Raw spectrum data were used to search against the UniProtKB/Swiss-Prot protein database for bovine. Main search parameters included: trypsin, maximum missed cleavage site of two, precursor mass tolerance of 1.5 Da, fragment mass tolerance of 0.8 Da, and variable modification of oxidation/hydroxylation of methionine, proline, and lysine (+15.995Da). Decoy database search was performed to calculate false discovery rate (FDR). Proteins containing one or more peptides with FDR≤0.05 were considered positively identified and reported. Total number of peptide spectral matches (PSMs) for each protein reported by Protein Discoverer 1.4 was used

for quantification.

### 3D-bioplotting of osteogenic and chondrogenic scaffold phases

We have previously optimized the geometry and assessed the biocompatibility of the 3D-bioplotted scaffolds used in this study(42, 43). Using the methods described previously, the base computer assisted design (CAD) models of the scaffolds were created in Solidworks (Dassault Systèmes SOLIDWORKS Corp., Waltham, MA) as discs (Ø14.5 x 2 mm) to fit the individual wells of a standard 24-well culture plate. The STL file, sliced into 6 layers, was positioned onto the bioplotting stage (BioplotterRP, EnvisionTEC GmbH, Gladbeck, Germany). To begin scaffold fabrication on the 3D-Bioplotter (EnvisionTEC GmbH), the strand-pore geometry previously determined to be favorable for hASC(42) and experimentally optimized process parameters were assigned (Table 1) in VisualMachines. 6.8 g of material (80% PCL-20% TCP composite for osteogenic scaffolds; 100% PCL for chondrogenic scaffolds) was pre-

heated in the high temperature cartridge and the scaffolds 3D-bioplotted, one at a time. Post-fabrication, all scaffolds were weighed (AR2140 digital balance, Ohaus Corp., Parsippany, NJ) and characterized for strand width and interstrand spacing (50X magnification, KH-7700 microscope, Hirox, Hackensack, NJ). From each scaffold, for each dimension metric, measurements were taken at 8 random locations. All scaffolds were sterilized for 30 minutes in 70% ethanol, and washed three times with sterile 1X PBS and once with complete growth medium (CGM, defined below).

To prepare the PCL-dECM scaffolds for chondrogenic analyses, 200 µL of the decellularized bovine cartilage ECM hydrogel (protocol described above) was added onto each sterilized 3D-bioplotted 100% PCL scaffold. The hydrogel-infused scaffolds were placed in an incubator (37°C, 5% CO<sub>2</sub>) overnight to allow the gel to solidify, and were then placed in 24-well plates prior to seeding with hASC.

#### Characterization of calcium release from PCL-TCP scaffolds

PCL-TCP scaffolds were placed into separate wells in a 24-well plate (n = 3 per time point) and treated for 30 minutes with 70% ethanol to mimic the sterilization process used in later cell culture experiments. Ethanol treatment also increased the hydrophilicity of the scaffold for the Ca<sup>2+</sup> release experiments. The scaffolds were then fully submerged in 0.5 mL of phosphate buffered saline (PBS) and maintained in an incubator (37°C, 5% CO<sub>2</sub>) for the duration of the study. PBS was collected at time points of 1, 2, 7, 14, and 28 days and stored at -25°C until assayed. A Calcium LiquiColor® Assay (StanBio, Boerne, TX) was used to quantify calcium content of each sample. Using the measured weight of the scaffolds along with the molar mass of calcium ions (Ca<sup>2+</sup> 40.08 g•mol<sup>-1</sup>), orthophosphates (PO<sub>4</sub><sup>3-</sup>94.97 g•mol<sup>-1</sup>), and TCP [Ca<sub>3</sub>(PO<sub>4</sub>)<sub>2</sub> 310.18 g•mol<sup>-1</sup>] the weight of Ca<sup>2+</sup> doped within each scaffold was calculated to be 11.9 µg.

# Human ASC isolation, expansion, and in vitro culture of scaffolds

Excess adipose tissue was collected from five female pre-menopausal donors (ages 24 to 36) in accordance with an approved IRB protocol at UNC Chapel Hill (IRB 04-1622), and hASC were isolated from the tissue as previously described by our lab and others(44, 45). Cells were expanded in CGM comprised of alpha-modified minimal essential medium (α-MEM with Lglutamine) (Invitrogen, Carlsbad CA), 10% fetal bovine serum (FBS) (Premium Select, Atlanta Biologicals, Lawrenceville GA), 200 mM L-glutamine, and 100 I.U. penicillin/ 100 μg/mL streptomycin (Mediatech, Herndon VA). Human ASC were cultured (37°C, 5% CO<sub>2</sub>) until reaching 80% confluency changing the media every 3 days, and then passaged using trypsin-EDTA (Invitrogen). A superlot was generated by pooling equal numbers of cells from the five individual donor cell lines into a single culture vessel as previously described by our lab, and characterized for multilineage differentiation potential by culturing for 14 days in osteogenic differentiation medium (Minimum Essential Medium, alpha modified supplemented with 10% FBS, 2 mM L-glutamine, 100 U/mL penicillin, 100 mg/mL streptomycin, 50 µM ascorbic acid, 0.1 μM dexamethasone, and 10mM β -qlycerolphosphate) and adipogenic differentiation medium (Minimum Essential Medium, alpha modified supplemented with 10% FBS, 2 mM L-glutamine, 100 U/mL penicillin, 100 mg/mL streptomycin,1 μM dexamethasone, 5 μg/mL insulin, 100 μΜ indomethacin and 500 µM isobutylmethylxanthine), ensuring the superlot differentiation was representative of an average of the five cell lines(45). All experiments were run using superlot passage 6 or lower.

21

22

23

24

1

2

3

4

5

6

7

8

9

10

11

12

13

14

15

16

17

18

19

20

To assess osteogenesis, PCL-TCP scaffolds (n = 8) were seeded with hASC at a density of 250,000 cells per scaffold and incubated for 28 days in CGM. To assess chondrogenesis, 250,000 hASC in 1mL of CGM were seeded onto each PCL-dECM scaffold (n = 9) and cultured for 28 days in CGM(25, 41).

26

### Osteogenic differentiation assays

Osteogenic differentiation was assessed via calcium staining and endogeonous alkaline phosphatase activity (ALP). Scaffolds were fixed for 30 minutes in formalin at day 28 and stained via Alizarin Red (Sigma-Aldrich) for calcium accretion. Endogenous ALP activity was assessed at days 14 and 28. Scaffolds were collected in 1 mL RIPA buffer (Fisher). ALP was quantified with the Alkaline Phosphatase Liquicolor Test (Stanbio, Boerne, TX), using the P-nitrophenylphosphate methodology, as described previously(46). ALP activity was normalized to total protein content quantified using a Pierce Micro BCA Protein Assay Kit (Thermo Fisher Scientific). Additionally, an Alamar Blue assay (AbD Serotec, Raleigh, NC) was used to assess metabolic activity of the seeded cells at days 14 and 28.

11

12

10

1

2

3

4

5

6

7

8

9

# Chondrogenic differentiation assays

- 13 Safranin-O staining
- 14 Safranin-O staining was used to determine proteoglycan content of the seeded scaffolds.
- 15 Scaffolds were fixed in 10% formalin and first stained with Weigert's Iron Hematoxylin stain
- 16 (Sigma-Aldrich) for 10 minutes, then rinsed under running tap water. They were then stained
- 17 with fast green for 5 minutes and washed in 1% acetic acid, followed by an 8 minute Safranin-O
- 18 stain.

19

- 20 Alcian Blue staining
- 21 Alcian Blue staining was used to determine sulfated glycosaminoglycan (sGAG) content of the
- 22 seeded scaffolds. After fixation as described above, scaffolds were stained in an Alcian blue
- 23 solution (Sigma-Aldrich) for 30 minutes, washed under running tap water for 2 minutes, and
- 24 rinsed with deionized water. Scaffolds were then stained with 0.1% Nuclear Fast Red (Sigma-
- 25 Aldrich) for 5 minutes, tap water for 1 minute.

#### Quantitative RT-PCR

1

2

3

4

5

6

7

8

9

10

11

12

13

14

15

16

17

18

19

20

21

22

23

24

25

26

Quantitative real-time RT-PCR was used to determine relative mRNA expression changes of chondrogenic-specific genes in seeded scaffolds with dECM or without dECM and osteogenicspecific genes seeded scaffolds with TCP or without TCP (n = 5 per condition). Total RNA was extracted directly from the scaffolds after 28 days in culture. following the TRIZOL protocol (Thermo Fisher Scientific). Briefly, the scaffolds were submerged in 1 mL of TRIZOL followed by a chloroform (Ricca Chemical Co. Arlington, TX) extraction. The precipitated solution was incubated with an equal volume of isopropanol (Sigma-Aldrich) and 1 µL glycoblue (Thermo Fisher Scientific) at room temperature for 10 minutes. The solution was centrifuged, and the precipitate was washed once with 75% ethanol (Koptec) and centrifuged again to collect the RNA containing pellet. The RNA was resuspended in DEPC treated water (Mediatech) and the yield was calculated using a NanoDrop 2000 Spectrophotometer (Sigma-Aldrich). To quantify mRNA expression, 120 ng of total RNA was reverse transcribed into cDNA using a high capacity first strand cDNA synthesis kit (OriGene, Rockville, MD) as per manufacturer's instructions, gRT-PCR was performed using 1 µL cDNA, 10 µL SYBR Green (Applied Biosystems, Foster City, CA), 7 µL DEPC treated water (Mediatech), and 1 µL forward and reverse primer (IDT DNA) for the following genes: b-actin '5-CACTCTTCCAGCCTTCCTTC-3', '3-TGTAGGCGTTTCTGGACATG-5', Osteopontin 5'-CTCAGGCCAGTTGCAGCC-3', 3'-CTCTTAACGTCACTAAACGAAAAC-5' **ALP** 5'-GACCCTTGACCCCCACAAT-3', 3'-TCCCCTGTACGTCATGCTCG-5' Collagen1a2 '5-AGAGTGGAGCAGTGGTTACTA-3', '3-GAGATGACCGCTTTGGACATAG-5', Aggrecan '5-CAGGCAGATCACTTGAGGTTAG-3'. '3-ACATTAGGGTCGATGATCCCTC-5', Sox9 '5-TGACCTATCCAAGCGCATTAC-3', **'3**-ATTTGGGAGAAGTCTCGTTCG-5'. qRT-PCR was performed using an ABI 7000 Sequence Detection system (Applied Biosystems). Samples were assayed in triplicate in one run (40 cycles) per gene. qRT-PCR

data were analyzed using the  $2^{-\Delta\Delta CT}$  method as described previously(47) with  $\beta$ -actin as the

1 housekeeping control. Relative quantification values are presented as fold changes in gene

expression relative to the control group.

3

2

# 4 Fabrication and in vitro hASC culture of integrated full thickness osteochondral scaffolds

- 5 Electrospun layer fabrication
- 6 Electrospun PCL matrices, designed to mimic the function of the tidemark layer, were
- 7 separately fabricated prior to 3D-bioplotting of the full thickness multiphasic scaffold. PCL was
- 8 dissolved in chloroform and dimethylformamide (3:1 ratio by volume) to create a 12% solution.
- 9 The solution was mixed continuously at 80°C for 4 hours. The PCL solution was electrospun
- immediately after preparation at a feed rate of 0.7 µL/hour using 15 kV and then cut into discs
- 11 (Ø14.5 mm X 30 μm) to match the 3D-bioplotted scaffolds.

12

13

- Full thickness scaffold assembly
- 14 The integrated triphasic scaffolds (Ø14.5 x 6 mm) were designed to resemble the overall
- 15 organization of the osteochondral complex, and the manufacturing method used has been
- described previously(42, 43). The material composition, strand-pore geometries, and process
- parameters for the bone (PCL-TCP), cartilage (PCL-dECM), and tidemark (electrospun PCL)
- 18 phases were as described above. To create the scaffolds in a single build, the PCL-TCP phase
- 19 (Ø14.5 x 4 mm, 12 layers) was bioplotted first. Immediately upon completion of the topmost
- 20 PCL-TCP layer, the process was briefly halted, and the pre-fabricated electrospun PCL disc
- 21 (Ø14.5 mm X 30 µm) was overlaid on top of the partial scaffold. 3D-bioplotting was then
- 22 continued with the appropriate height offset adjustment to create the PCL phase (Ø14.5 x 2 mm,
- 23 6 layers).

- 25 A two-day hASC seeding protocol was used to seed the full thickness scaffolds (n = 3). On the
- 26 first day, the scaffolds were sterilized with successive 70% ethanol and PBS washes. The

chondrogenic phase was prepared by submersing the 3D-bioplotted PCL in neutralized dECM solution and incubating at 37°C for 1 hour. After the hydrogel had solidified, the scaffold was flipped chondrogenic PCL phase up, seeded with 250,000 hASC, and incubated overnight (37°C, 5% CO<sub>2</sub>). On the second day, the scaffolds were flipped PCL-TCP phase up, and seeded with an additional 500,000 hASC. The total number of cells used in these scaffolds was tripled in the full osteochondral scaffolds because they were three times thicker than the individual chondrogenic and osteogenic scaffolds. Each osteochondral scaffold was incubated in 3 mL CGM for 28 days. Media was changed and scaffolds were flipped every two days.

#### Characterization of full thickness osteochondral scaffolds

After 28 days of culture, full thickness scaffolds (n = 3 unseeded, n = 3 seeded) were fixed for 30 minutes in formalin. Total scaffold fixation and staining has been utilized previously to demonstrate calcium distribution in bone scaffolds (48, 49). Scaffolds were sectioned with a scalpel into 1 mm thick cross-sections for histological analyses. Sections were stained via Alizarin Red, Alcian Blue, and Safranin-O/fast green, as described above.

### Characterization of the effects of electrospun PCL tidemark matrix

PCL scaffolds were fabricated with (n = 4) or without (n = 4) an electrospun PCL tidemark matrix to evaluate cell migration characteristics. Scaffolds had an overall thickness of 6mm, and tidemarks were added to split the scaffolds into 2 mm and 4 mm sections. Each scaffold was seeded by suspending 250,000 cells in 150 µL of media in a well of a 24-well plate for 4 hours with the 2 mm scaffold section facing downwards into the cell suspension. After 4 hours, the 2 mm section was flipped upwards in a new 24-well plate and 1 mL of media was added to the scaffold. This yielded a scaffold with the top 2 mm section seeded with cells but the bottom 4 mm section remaining unseeded. Sterile, non-tissue culture treated plates were used to prevent cells from attaching to the plate. Half media changes were carried out every other day for 28

- days. After 28 days, scaffolds were fixed and stained via Hematoxylin and Eosin (H&E) for cell
- 2 nuclei and protein distribution. The scaffolds were submerged in Hematoxylin for 10 minutes,
- 3 washed under running tap water for 5 minutes, rinsed in deionized water, stained via Eosin for 2
- 4 minutes, rinsed in deionized water and immediately imaged on an EZ4D histology microscope
- 5 (Leica, Wetzlar, Germany).

6

7

# Statistical Analyses

- 8 Statistical analyses were performed using Prism (version 5.0a, GraphPad Software). Student t-
- 9 test was used to assess significant differences between two groups, and ANOVA with Tukey
- 10 post-hoc test was used to asses significant differences between 3 or more groups. All Graphs
- 11 are shown in the form of mean  $\pm$  SEM.

12

13

#### RESULTS

- 14 PCL-TCP scaffolds induced osteogenic differentiation of hASC
- 15 Controlled release assays of calcium demonstrated that calcium was released from the PCL-
- 16 TCP composite scaffolds throughout the 28 day incubation period, with a large initial release
- within the first 24 hours (**Figure 2A**).

- 19 To determine how the calcium released from the scaffolds affected the seeded hASC, assays
- were performed to assess osteogenic differentiation. Enzymatic ALP activity, a hallmark of
- osteogenic differentiation(46), was elevated PCL-TCP composition scaffolds when compared to
- 22 pure PCL scaffolds (**Figure 2B**). This trend was observed at both the 14 (p = 0.115) and 28-day
- 23 (p = 0.073) time points. At the gene expression level, ALP mRNA expression was elevated in
- 24 PCL-TCP scaffolds at day 14 (p = 0.323), but was reduced relative to the control by day 28 (\*p =
- 25 0.018) (Figure 2C). Osteopontin was elevated in PCL-TCP at day 14 (\*p < 0.001) (Figure 2D).
- 26 End product calcium staining demonstrated that hASC cultured on PCL-TCP scaffolds accreted

1 calcium while hASC cultured on PCL scaffolds alone did not accrete calcium. Acellular PCL-

TCP controls showed mild Alizarin Red staining due to the TCP found within the scaffold, but

seeded PCL-TCP stained dark red throughout the scaffold, indicating hASC-mediated calcium

accretion (Figure 2E). Scaffolds without TCP showed no Alizarin Red staining.

5

6

7

8

9

10

11

12

2

3

4

We have previously demonstrated that bioplotted PCL scaffolds support viability and

proliferation of hASC(42, 43). To assess the cell viability and proliferation within the current

study, Alamar Blue was used to detect metabolic activity of the seeded cells (Figure 2F). In this

assay, a higher percent reduction corresponds to higher metabolic activity of the assayed cells.

On both pure PCL scaffolds and PCL-TCP composite scaffolds, metabolic activity increased

between days 14 and 28, which suggests continued proliferation of the cells throughout the 4-

week culture period. Metabolic activity was higher in PCL-TCP composite scaffolds than in pure

13 PCL scaffolds at both time points.

14

15

17

18

19

20

#### PCL-dECM scaffolds induced chondrogenic differentiation of hASC

16 The dECM hydrogel used to induce hASC chondrogenic differentiation was analyzed using

proteomic analysis to ensure that the primary ECM proteins were preserved after the

decellularization process(50) (Figure 3). We identified 775 unique peptides corresponding to

proteoglycans, collagens and other extracellular matrix proteins. Figure 3 shows the relative

abundances of the most prevalent proteins.

21

22

23

24

25

26

Human ASC were then seeded on 3D-bioplotted scaffolds with or without dECM and evaluated

for chondrogenic differentiation. Histological staining showed more robust proteoglycan

(Safranin-O) and sulfated glycosaminoglycan (Alcian Blue) staining in scaffolds that were

seeded when compared to their unseeded controls, even in the absence of dECM (Figure 4).

Unseeded scaffolds without dECM did not stain for either proteoglycan or sulfated

glycosaminoglycan content, as expected. Moderate matrix deposition was observed on the hASC seeded scaffolds without dECM, while the unseeded scaffolds with a dECM coating stained positive for proteoglycan or sulfated glycosaminoglycan from the dECM itself. The unseeded dECM had regions of degradation and lacked organization. Seeded scaffolds with dECM coating had an evenly distributed matrix that had undergone remodeling from the cells. In addition, qualitative observation of the scaffolds showed that more matrix was deposited on the dECM seeded scaffolds than the unseeded scaffolds, indicating end product accretion of chondrogenic differentiation. Human ASC chondrogenesis was further evaluated by qPCR with similar results (Figure 4M-O). Although no statistical significance was observed in this data set the following trends were observed. Seeded PCL-dECM scaffolds exhibited increased expression of sox9, an early marker of chondrogenic differentiation(51). After 28 days in culture, aggrecan expression in PCL-dECM scaffolds was elevated. Furthermore, Collagen I expression in dECM coated scaffolds was reduced relative to hASC seeded on non-dECM scaffolds at day 14 but elevated at day 28.

### Full thickness osteochondral scaffolds exhibited site-specific tissue differentiation

Histological staining of the full-thickness scaffolds revealed site-specific osteochondral tissue characteristics (Figure 5). The PCL-TCP region of the scaffold stained dark red with the Alizarin Red staining, indicating calcium accretion within this region of the scaffold. The PCL-dECM portion of the scaffold stained with Safranin-O indicated the presence of proteoglycans, and Alcian Blue indicated the presence of sulfated glycosaminoglycans. Alizarin Red, Alcian Blue, and Safranin-O staining was more robust in the hASC seeded scaffolds than the non-seeded controls, demonstrating that end-product expression rather than the biomimetic nature of the scaffolds themselves contributed to at least part of the observed staining patterns. In addition, the bone-like portion of the scaffold and cartilage-like portion of the scaffold transitioned cleanly from one phase to the other at the electrospun tide mark.

# The electrospun tidemark prevents cell migration

H&E staining of scaffolds that were cultured with and without an electrospun layer to mimic a osteochondral tidemark displayed different cell distributions (**Figure 6**). Scaffolds that included a tidemark displayed much more robust nuclear and protein staining on the seeded side of the scaffolds and reduced cell nuclear and protein staining on the non-seeded side. However, scaffolds that did not include a tidemark had much more homogenous cell distribution throughout the scaffold. The scaffolds without an electrospun layer had a greater density of nuclei on the non-seeded side of the scaffold than the scaffolds that contained an electrospun layer.

### DISCUSSION

The objective of this study was to develop a tissue engineered osteochondral construct that closely resembled the native architecture of osteochondral tissue using a single cell source of abundant and accessible hASC. A triphasic scaffold was created by combining two commonly used techniques (electrospinning and 3D printing), and incorporating chemical cues within the different layers of the scaffold to induce site-specific osteogenic and chondrogenic differentiation of the hASC. Specifically, TCP was used to induce osteogenic differentiation and dECM was used to induce chondrogenic differentiation.

TCP has previously been shown to drive osteogenic differentiation (25, 52, 53). It has been demonstrated to induce differentiation in both 2D culture and 3D electrospun constructs (52). More recently we demonstrated that polylactic acid nanofibrous scaffolds containing 20% TCP nanoparticles drove osteogenic differentiation of hASC and that elevated calcium inhibited chondrogenic differentiation, even in the presence of chondrogenic growth factors (25). Here we confirm that scaffolds containing 20% TCP drive cells to differentiate down the osteogenic

lineage by demonstrating that osteogenic markers are upregulated and end product calcium accretion is enhanced in the presence of TCP. When compared to previously reported electrospun scaffolds containing TCP, calcium is released at a much slower rate from the 3D-bioplotted PCL-TCP scaffolds. Specifically, we have previously demonstrated a burst release of 50% calcium within the first 24 hour in electrospun scaffolds (53). Here, we only witnessed a release of 0.54 μg (4.5% of total Ca<sup>2+</sup>) with the first 24 hours, and a sustained release over the course of 28 days. Even after 28 days, only 1.04 μg (8.7% of total Ca<sup>2+</sup>) had been released from the scaffolds, suggesting that calcium-based signaling would be sustained long term in hASC seeded upon these scaffolds, which could be favorable for implantation of the tissue engineered constructs *in vivo*.

The cartilage layer of the 3D-bioplotted scaffold was coated with cartilage dECM in order to drive chondrogenic differentiation of the hASC without the need of growth factor supplementation. In recent years, there has been great interest in the use of dECM for osteochondral tissue engineering(54). Since naturally derived ECM has a complex array of bioactive molecules, it is ideal to signal cells to adhere(55), proliferate(56), and differentiate (57, 58). In cartilage tissue engineering, the use of a bioactive biomaterial is especially useful since cartilage is avascular and cannot easily circulate growth factors and nutrients(54). Previously, biphasic osteochondral scaffolds have been produced via combination of decellularized cartilage and decellularized bone(59). In addition, monophasic decellularized equine cartilage matrix implants have been used to generate biphasic osteochondral regeneration, although potential vascularization and ossification of the neo-cartilage is a concern(54). In this study, we demonstrate that most of the crucial extracellular proteins of articular cartilage are preserved after the decellularization step. We used this naturally-derived hydrogel to direct chondrogenic differentiation of hASC, even in the absence of supplemental chondrogenic growth factors. This is in accordance with previous literature, that demonstrated bone marrow derived mesencyhmal

stem cells can be shunted down the chondrogenic lineage via stimulation with cartilage fragments without the use of growth factors(60). Additionally, in this study, we demonstrate that hASC have a synergistic effect with the dECM. We observe that seeded dECM displayed more robust proteoglycan and sulfated glycosaminoglycan than the non-seeded dECM. Upon closer examination of the histology, the seeded dECM appeared to have been remodeled and augmented by the hASC whereas the non-seeded dECM suffered from degradation. This indicates that while the dECM is driving chondrogenic differentiation, the cells are simultaneously modifying the structural properties of the dECM. The results from both dECM and TCP studies drove the development of single osteochondral scaffolds, which can be cultured in complete growth medium alone (without any additional osteogenic and/or chondrogenic growth factors) to induce both hASC chondrogenesis and osteogenesis in desired site-specific locations.

A novel aspect of our triphasic design was the incorporation of a thin eletropsun PCL layer between the osteogenic and chondrogenic layers to mimic the natural tidemark found in osteochondral tissue. When engineering osteochondral implants, it is a concern that the neocartilage would ossify if implanted *in vivo* long term(54). The incorporation of the electrospun tidemark layer is important because it prevents cell migration between the cartilage and bone layers. The *in vitro* results of this study indicate that hASC migration is inhibited in scaffolds that include an electrospun layer. This layer could potentially act as a barrier to prevent blood vessel invasion into the cartilage layer when implanted *in vivo*.

Since the overall objective of this study was to generate an osteochondral implant with sitespecific characteristics, the global protein and calcium distributions within the 3D multi-phasic scaffolds were assessed. The scaffolds were cultured for 4 weeks, which gave the bioactive factors and cells seeded upon those layers ample time to interact with each other. Since both the osteogenic side and chondrogenic side were cultured simultaneously within the same dish, interactions between the two phases cannot be avoided, and we did not attempt to mitigate these interactions. Since bone and cartilage are known to crosstalk extensively *in vivo* (61), this mimics native osteochondral signaling. Staining revealed that site-specific characteristics within the 3D architecture are maintained throughout the duration of the culture, despite these interactions.

Although the current study presents a novel system for producing site specific differentiation of hASC within a single scaffold, a limitation of the current study is that only *in vitro* characterization was performed. While *in vitro* characterization is an essential step before moving into animal models, this scaffold system will require *in vivo* validation in future studies. Specifically, in order to assess the potential for large scale preclinical studies and eventual clinical translation, this scaffold system must be validated in a large animal model to ensure that the scaffold does not elicit an immune response and achieves the desired healing *in vivo* (62). Additionally, for this study we use decellularized porcine ECM as it is easily obtainable and is amenable to future porcine in vivo studies. However, for clinical purposes, large amounts of human cartilage will be more difficult to obtain, and therefore an immune response to porcine tissue will have to be considered. Although the tissue is decellularized and sterilized, further studies to assess cross-species immune response will be necessary prior to clinical studies.

This study is the first to incorporate TCP and dECM in a single 3D-bioplotted scaffold to induce site specific differentiation of hASC as a single cell source to generate a full osteochondral construct. Recent research in osteochondral scaffold fabrication has utilized 3D printing approaches to generate patient specific scaffolds in a precise and highly-tunable manner. In this study we take advantage of recent advances in biofabrication to generate scaffolds that are thick enough to repair full-thickness osteochondral defects, unlike electrospun scaffolds. We

show, for the first time, that the incorporation of an electrospun layer can inhibit cell migration between scaffold layers, which may be highly advantageous *in vivo* by preventing blood vessels from invading the chondrogenic portion of the scaffold. The scaffold design described in this study minimizes rejection by using an abundant and accessible source of autologous stem cells, and our biofabrication techniques allow for a precise, customizable methodology to rapidly produce the scaffold. With further development, this novel approach could hold great potential to treat OA patients by offering a patient-specific, less-invasive alternative to full joint replacement.

### **ACKNOWLEDGEMENTS**

Special thanks to Saahil Mehendale for support on scaffold design, Sonya Sonnenberg for assistance with decellularization of articular cartilage and Alison Fragale for illustrating Figure 1. This work was supported in part by the OREF Research Grant in Stem Cells and Regenerative Medicine Grant # 15-059 (JS, EGL, RAS), National Space Biomedical Research Institute through NASA NCC9-58 (LFM), NIH/NIBIB 1R03EB008790 (EGL), NIH/CTSA 550KR71418 (EGL), NIH/CTSA 550KR61325 (EGL), NSF/CBET 1133427 and 1702841 (EGL) and NIH/COBRE P20GM109095 (JTO) and NIH/INBRE P20GM103408 (JTO). The 3D-bioplotting aspect of this study was supported by a grant from NCSU Faculty Research and Professional Development (FRPD) program and NSF/CBET 1703466 (RAS).

### REFERENCES

1

2 References

- 1. Kotlarz, H, Gunnarsson, CL, Fang, H, Rizzo, JA. Insurer and out-of-pocket costs of osteoarthritis in the US: evidence from national survey data. Arthritis Rheum 2009;60:3546-3553.
- the US: evidence from national survey data. Arthritis Rheum 2009;60:3546-3553.
   Revell, CM, Athanasiou, KA. Success rates and immunologic responses of autogenic, allogenic, and xenogenic treatments to repair articular cartilage defects. Tissue Eng Part B Rev 2009;15:1-15.
- 7 3. Niemeyer, P, Pestka, JM, Kreuz, PC, Erggelet, C, Schmal, H, Suedkamp, NP, Steinwachs, M.
  8 Characteristic complications after autologous chondrocyte implantation for cartilage defects of the knee joint. Am J Sports Med 2008;36:2091-2099.
- 4. Yousefi, AM, Hoque, ME, Prasad, RG, Uth, N. Current strategies in multiphasic scaffold design for osteochondral tissue engineering: A review. J Biomed Mater Res A 2015;103:2460-2481.
- 5. Gilbert, JE. Current treatment options for the restoration of articular cartilage. Am J Knee Surg 1998;11:42-46.
- 6. Getgood, A, Brooks, R, Fortier, L, Rushton, N. Articular cartilage tissue engineering: today's research, tomorrow's practice? J Bone Joint Surg Br 2009;91:565-576.
- Atesok, K, Doral, MN, Karlsson, J, Egol, KA, Jazrawi, LM, Coelho, PG, Martinez, A, Matsumoto,
   T, Owens, BD, Ochi, M, Hurwitz, SR, Atala, A, Fu, FH, Lu, HH, Rodeo, SA. Multilayer scaffolds in orthopaedic tissue engineering. Knee Surg Sports Traumatol Arthrosc 2014.
- Mobasheri, A, Csaki, C, Clutterbuck, AL, Rahmanzadeh, M, Shakibaei, M. Mesenchymal stem cells in connective tissue engineering and regenerative medicine: applications in cartilage repair and osteoarthritis therapy. Histol Histopathol 2009;24:347-366.
- Valonen, PK, Moutos, FT, Kusanagi, A, Moretti, MG, Diekman, BO, Welter, JF, Caplan, AI,
   Guilak, F, Freed, LE. In vitro generation of mechanically functional cartilage grafts based on adult
   human stem cells and 3D-woven poly(epsilon-caprolactone) scaffolds. Biomaterials 2010;31:2193 2200.
- Huang, AH, Farrell, MJ, Mauck, RL. Mechanics and mechanobiology of mesenchymal stem cell-based engineered cartilage. J Biomech 2010;43:128-136.
- Huang, CY, Reuben, PM, D'Ippolito, G, Schiller, PC, Cheung, HS. Chondrogenesis of human bone marrow-derived mesenchymal stem cells in agarose culture. Anat Rec A Discov Mol Cell Evol Biol 2004;278:428-436.
- Lee, HJ, Yu, C, Chansakul, T, Hwang, NS, Varghese, S, Yu, SM, Elisseeff, JH. Enhanced
   chondrogenesis of mesenchymal stem cells in collagen mimetic peptide-mediated
   microenvironment. Tissue Eng Part A 2008;14:1843-1851.
- van Osch, GJ, Brittberg, M, Dennis, JE, Bastiaansen-Jenniskens, YM, Erben, RG, Konttinen, YT,
   Luyten, FP. Cartilage repair: past and future--lessons for regenerative medicine. J Cell Mol Med
   2009;13:792-810.
- Huang, AH, Stein, A, Mauck, RL. Evaluation of the complex transcriptional topography of mesenchymal stem cell chondrogenesis for cartilage tissue engineering. Tissue Eng Part A 2010;16:2699-2708.
- Huang, AH, Farrell, MJ, Kim, M, Mauck, RL. Long-term dynamic loading improves the mechanical properties of chondrogenic mesenchymal stem cell-laden hydrogel. Eur Cell Mater 2010;19:72-85.
- Hillel, AT, Taube, JM, Cornish, TC, Sharma, B, Halushka, M, McCarthy, EF, Hutchins, GM, Elisseeff, JH. Characterization of human mesenchymal stem cell-engineered cartilage: analysis of its ultrastructure, cell density and chondrocyte phenotype compared to native adult and fetal cartilage. Cells Tissues Organs 2010;191:12-20.
- 47 17. Deans, TL, Elisseeff, JH. Stem cells in musculoskeletal engineered tissue. Curr Opin Biotechnol 2009;20:537-544.

- Sundelacruz, S, Kaplan, DL. Stem cell- and scaffold-based tissue engineering approaches to osteochondral regenerative medicine. Semin Cell Dev Biol 2009;20:646-655.
   Swieszkowski, W, Tuan, BH, Kurzydlowski, KJ, Hutmacher, DW. Repair and regeneration of the company o
- 3 19. Swieszkowski, W, Tuan, BH, Kurzydlowski, KJ, Hutmacher, DW. Repair and regeneration of osteochondral defects in the articular joints. Biomol Eng 2007;24:489-495.
- Grayson, WL, Bhumiratana, S, Grace Chao, PH, Hung, CT, Vunjak-Novakovic, G. Spatial regulation of human mesenchymal stem cell differentiation in engineered osteochondral constructs: effects of pre-differentiation, soluble factors and medium perfusion. Osteoarthritis Cartilage 2010;18:714-723.
- 9 21. Hung, CT, Lima, EG, Mauck, RL, Takai, E, LeRoux, MA, Lu, HH, Stark, RG, Guo, XE, Ateshian, GA. Anatomically shaped osteochondral constructs for articular cartilage repair. J Biomech 2003;36:1853-1864.
- 12 22. Lima, EG, Mauck, RL, Han, SH, Park, S, Ng, KW, Ateshian, GA, Hung, CT. Functional tissue engineering of chondral and osteochondral constructs. Biorheology 2004;41:577-590.
- 23. Schaefer, D, Martin, I, Shastri, P, Padera, RF, Langer, R, Freed, LE, Vunjak-Novakovic, G. In vitro generation of osteochondral composites. Biomaterials 2000;21:2599-2606.
- Shimomura, K, Moriguchi, Y, Murawski, CD, Yoshikawa, H, Nakamura, N. Osteochondral tissue engineering with biphasic scaffold: current strategies and techniques. Tissue Eng Part B Rev 2014;20:468-476.
- Mellor, LF, Mohiti-Asli, M, Williams, J, Kannan, A, Dent, MR, Guilak, F, Loboa, EG.
   Extracellular Calcium Modulates Chondrogenic and Osteogenic Differentiation of Human
   Adipose-Derived Stem Cells: A Novel Approach for Osteochondral Tissue Engineering Using a
   Single Stem Cell Source. Tissue Eng Part A 2015;21:2323-2333.
- 23 26. Tampieri, A, Sandri, M, Landi, E, Pressato, D, Francioli, S, Quarto, R, Martin, I. Design of graded biomimetic osteochondral composite scaffolds. Biomaterials 2008;29:3539-3546.
- 25 27. Jiang, J, Tang, A, Ateshian, GA, Guo, XE, Hung, CT, Lu, HH. Bioactive stratified polymer
   26 ceramic-hydrogel scaffold for integrative osteochondral repair. Ann Biomed Eng 2010;38:2183 27 2196.
- 28. Kon, E, Mutini, A, Arcangeli, E, Delcogliano, M, Filardo, G, Nicoli Aldini, N, Pressato, D, Quarto,
   R, Zaffagnini, S, Marcacci, M. Novel nanostructured scaffold for osteochondral regeneration: pilot
   study in horses. J Tissue Eng Regen Med 2010;4:300-308.
- 31 29. Kon, E, Delcogliano, M, Filardo, G, Pressato, D, Busacca, M, Grigolo, B, Desando, G, Marcacci,
   32 M. A novel nano-composite multi-layered biomaterial for treatment of osteochondral lesions:
   33 technique note and an early stability pilot clinical trial. Injury 2010;41:693-701.
- 30. Kon, E, Delcogliano, M, Filardo, G, Fini, M, Giavaresi, G, Francioli, S, Martin, I, Pressato, D, Arcangeli, E, Quarto, R, Sandri, M, Marcacci, M. Orderly osteochondral regeneration in a sheep model using a novel nano-composite multilayered biomaterial. J Orthop Res 2010;28:116-124.
- 37 31. Erisken, C, Kalyon, DM, Wang, H. Viscoelastic and biomechanical properties of osteochondral tissue constructs generated from graded polycaprolactone and beta-tricalcium phosphate composites. J Biomech Eng 2010;132:091013.
- 40 32. Lu, HH, Subramony, SD, Boushell, MK, Zhang, X. Tissue engineering strategies for the regeneration of orthopedic interfaces. Ann Biomed Eng 2010;38:2142-2154.
- 42 33. Nukavarapu, SP, Dorcemus, DL. Osteochondral tissue engineering: current strategies and challenges. Biotechnol Adv 2013;31:706-721.
- 44 34. Pittenger, MF, Mackay, AM, Beck, SC, Jaiswal, RK, Douglas, R, Mosca, JD, Moorman, MA, Simonetti, DW, Craig, S, Marshak, DR. Multilineage potential of adult human mesenchymal stem cells. Science 1999;284:143-147.
- 47 35. Nordberg, RC, Loboa, EG. Our Fat Future: Translating Adipose Stem Cell Therapy. Stem Cells Transl Med 2015;4:974-979.
- 49 36. Awad, HA, Wickham, MQ, Leddy, HA, Gimble, JM, Guilak, F. Chondrogenic differentiation of adipose-derived adult stem cells in agarose, alginate, and gelatin scaffolds. Biomaterials 2004;25:3211-3222.

- Huang, JI, Zuk, PA, Jones, NF, Zhu, M, Lorenz, HP, Hedrick, MH, Benhaim, P. Chondrogenic potential of multipotential cells from human adipose tissue. Plast Reconstr Surg 2004;113:585-594.
   Zuk, PA, Zhu, M, Mizuno, H, Huang, J, Futrell, JW, Katz, AJ, Benhaim, P, Lorenz, HP, Hedrick,
- 3 38. Zuk, PA, Zhu, M, Mizuno, H, Huang, J, Futrell, JW, Katz, AJ, Benhaim, P, Lorenz, HP, Hedrick,
   4 MH. Multilineage cells from human adipose tissue: implications for cell-based therapies. Tissue
   5 Eng 2001;7:211-228.
   6 39. Huang, JI, Kazmi, N, Durbhakula, MM, Hering, TM, Yoo, JU, Johnstone, B. Chondrogenic
- Huang, JI, Kazmi, N, Durbhakula, MM, Hering, TM, Yoo, JU, Johnstone, B. Chondrogenic potential of progenitor cells derived from human bone marrow and adipose tissue: a patient-matched comparison. J Orthop Res 2005;23:1383-1389.
- Mehlhorn, AT, Niemeyer, P, Kaiser, S, Finkenzeller, G, Stark, GB, Sudkamp, NP, Schmal, H.
   Differential expression pattern of extracellular matrix molecules during chondrogenesis of
   mesenchymal stem cells from bone marrow and adipose tissue. Tissue Eng 2006;12:2853-2862.
- 12 41. Puetzer, JL, Petitte, JN, Loboa, EG. Comparative review of growth factors for induction of threedimensional in vitro chondrogenesis in human mesenchymal stem cells isolated from bone marrow and adipose tissue. Tissue Eng Part B Rev 2010;16:435-444.
- 42. Mehendale, S, Mellor, L, Taylor, M, Loboa, E, Shirwaiker, R. Characterizing the effects of 3D-bioplotted polycaprolactone (PCL) scaffold geometry on human adipose-derived stem cell viability and proliferation. Rapid Prototyping Journal 2017.
- Mellor, LF, Huebner, P, Cai, S, Mohiti-Asli, M, Taylor, MA, Spang, J, Shirwaiker, RA, Loboa, EG.
   Fabrication and Evaluation of Electrospun, 3D-Bioplotted, and Combination of Electrospun/3D-Bioplotted Scaffolds for Tissue Engineering Applications. Biomed Res Int 2017;2017:6956794.
- 44. Bernacki, SH, Wall, ME, Loboa, EG. Isolation of human mesenchymal stem cells from bone and
   adipose tissue. Methods Cell Biol 2008;86:257-278.
- 45. Bodle, JC, Teeter, SD, Hluck, BH, Hardin, JW, Bernacki, SH, Loboa, EG. Age-Related Effects on
   the Potency of Human Adipose-Derived Stem Cells: Creation and Evaluation of Superlots and
   Implications for Musculoskeletal Tissue Engineering Applications. Tissue Eng Part C Methods
   2014;20:972-983.
- 27 46. Bodle, JC, Rubenstein, CD, Phillips, ME, Bernacki, SH, Qi, J, Banes, AJ, Loboa, EG. Primary cilia: the chemical antenna regulating human adipose-derived stem cell osteogenesis. PLoS One 2013;8:e62554.
- 47. Livak, KJ, Schmittgen, TD. Analysis of relative gene expression data using real-time quantitative PCR and the 2(-Delta Delta C(T)) Method. Methods 2001;25:402-408.
- 32 48. Ma, R, Tang, S, Tan, H, Lin, W, Wang, Y, Wei, J, Zhao, L, Tang, T. Preparation, characterization, and in vitro osteoblast functions of a nano-hydroxyapatite/polyetheretherketone biocomposite as orthopedic implant material. Int J Nanomedicine 2014;9:3949-3961.
- 49. Lu, Y, Zhang, S, Liu, X, Ye, S, Zhou, X, Huang, Q, Ren, L. Silk/agarose scaffolds with tunable properties via SDS assisted rapid gelation. RSC Adv 2017.
- 50. Pu, X, Oxford, JT. Proteomic Analysis of Engineered Cartilage. Methods Mol Biol 2015;1340:263-278.
- 51. Kawakami, Y, Rodriguez-Leon, J, Izpisua Belmonte, JC. The role of TGFbetas and Sox9 during limb chondrogenesis. Curr Opin Cell Biol 2006;18:723-729.
- 41 52. McCullen, SD, Zhan, J, Onorato, ML, Bernacki, SH, Loboa, EG. Effect of varied ionic calcium on human adipose-derived stem cell mineralization. Tissue Eng Part A 2010;16:1971-1981.
- 43 53. Asli, MM, Pourdeyhimi, B, Loboa, EG. Release profiles of tricalcium phosphate nanoparticles
  44 from poly(L-lactic acid) electrospun scaffolds with single component, core-sheath, or porous fiber
  45 morphologies: effects on hASC viability and osteogenic differentiation. Macromol Biosci
  46 2012;12:893-900.
- 47 54. Benders, KE, van Weeren, PR, Badylak, SF, Saris, DB, Dhert, WJ, Malda, J. Extracellular matrix scaffolds for cartilage and bone regeneration. Trends Biotechnol 2013;31:169-176.
- Tavassoli, A, Matin, MM, Niaki, MA, Mahdavi-Shahri, N, Shahabipour, F. Mesenchymal stem cells can survive on the extracellular matrix-derived decellularized bovine articular cartilage scaffold. Iran J Basic Med Sci 2015;18:1221-1227.

- Vorotnikova, E, McIntosh, D, Dewilde, A, Zhang, J, Reing, JE, Zhang, L, Cordero, K, Bedelbaeva, K, Gourevitch, D, Heber-Katz, E, Badylak, SF, Braunhut, SJ. Extracellular matrix-derived products modulate endothelial and progenitor cell migration and proliferation in vitro and stimulate regenerative healing in vivo. Matrix Biol 2010;29:690-700.
- regenerative healing in vivo. Matrix Biol 2010;29:690-700.

  Gong, J, Sagiv, O, Cai, H, Tsang, SH, Del Priore, LV. Effects of extracellular matrix and neighboring cells on induction of human embryonic stem cells into retinal or retinal pigment epithelial progenitors. Exp Eye Res 2008;86:957-965.
- 8 58. Sellaro, TL, Ravindra, AK, Stolz, DB, Badylak, SF. Maintenance of hepatic sinusoidal endothelial cell phenotype in vitro using organ-specific extracellular matrix scaffolds. Tissue Eng 2007;13:2301-2310.
- Yang, Q, Peng, J, Lu, SB, Guo, QY, Zhao, B, Zhang, L, Wang, AY, Xu, WJ, Xia, Q, Ma, XL, Hu,
   YC, Xu, BS. Evaluation of an extracellular matrix-derived acellular biphasic scaffold/cell construct
   in the repair of a large articular high-load-bearing osteochondral defect in a canine model. Chin
   Med J (Engl) 2011;124:3930-3938.
- 15 60. Chen, CC, Liao, CH, Wang, YH, Hsu, YM, Huang, SH, Chang, CH, Fang, HW. Cartilage 16 fragments from osteoarthritic knee promote chondrogenesis of mesenchymal stem cells without 17 exogenous growth factor induction. J Orthop Res 2012;30:393-400.
- 18 61. Funck-Brentano, T, Cohen-Solal, M. Crosstalk between cartilage and bone: when bone cytokines matter. Cytokine Growth Factor Rev 2011;22:91-97.
- Cook, JL, Hung, CT, Kuroki, K, Stoker, AM, Cook, CR, Pfeiffer, FM, Sherman, SL, Stannard, JP.
   Animal models of cartilage repair. Bone Joint Res 2014;3:89-94.

Figure 1: Scaffold design for osteochondral tissue generation using a single cell source. In order to engineer a full osteochondral tissue using a single cell source, site-specific chemical cues were incorporated within the scaffold to induce site-specific differentiation of hASC. To achieve chondrogenesis, hASC were seeded within a 3D bioplotted PCL scaffold containing dECM. An electrospun layer of PCL was created and used to recreate the tidemark. To achieve osteogenesis, hASC were seeded within a 3D bioplotted scaffold comprised of 80% PCL and 20% TCP. Full composite scaffold shown on top right (scale bar = 5 mm).

Figure 2: Osteogenesis of hASC on 3D-bioplotted PCL-TCP scaffolds. A. The calcium release profile of osteogenic scaffolds (20% TCP 80% PCL) evaluated over 28 days. (bars represent standard error of the mean; different letters indicate statistical difference (p<0.05)). B. Endogenous ALP activity of seeded scaffolds at both the 14 and 28 day time points. Gene expression of C. ALP and D. Osteopontin. E. Alizarin Red staining indicated calcium accretion (scale bars = 200  $\mu$ m) F. Alamar Blue data indicating metabolic activity at days 14 and 28 of culture (\*\*\*p<0.001, \*p<0.05).

Figure 3: Proteomic analysis of dECM. Proteomic analysis of the dECM identified several cartilage proteins based on 775 unique peptides from articular cartilage. Proteoglycans: The most prevalent include fibromodulin, biglycan, PRELP, chondroadherin, decorin, aggrecan, link protein, lumican, and perlecan. Collagens: The most prevalent identified include COL2A1, COL6A3, COL9A1, COL12A1, COL6A2, COL6A1, COL15A1, COL11A2, and COL11A1. Noncollagenous ECM proteins: The most prevalent include fibronectin, Cartilage Oligomeric Matrix Protein (COMP), matrilin 3 (MATN3), cartilage intermediate layer protein 2 (CILP2), tenascin C (TNC), thrombospondin type 1 (THBS1), thrombospondin type 4 (THBS4), and clusterin.

- 1 Figure 4: Chondrogenic evaluation of 3D-bioplotted PCL-dECM scaffolds. A-D. Unstained
- 2 scaffolds of each condition E-H. Scaffolds stained for proteoglycan content via Safranin-O. I-L.
- 3 Scaffolds stained for sulfated glycosaminoglycans content via Alcian Blue (scale bars = 200
- 4 μm). Gene expression of M. sox9 and N. aggrecan, and O. collagen I.

5

- 6 Figure 5: Site-specific matrix distribution in full-thickness osteochondral scaffolds. A-B. Alizarin
- 7 red shows calcium deposition in the osteo layer, while C-D. Alcian Blue and E-F. Safranin-O
- 8 staining demonstrate increased proteoglycan and sulfated glycosaminoglycans in the superficial
- 9 cartilage layer (scale bars = 2 mm).

10

11

- Figure 6: Gross comparison of scaffolds stained via H&E shows that A. scaffolds without an
- 12 electrospun tidemark have relatively homogenous cellular distribution. B. Scaffolds that contain
- 13 an electrospun tidemark have more robust staining on the seeded side when compared to the
- 14 non-seeded side of the scaffold. Higher magnification of the far non-seeded sides of the
- scaffolds demonstrates that more nuclei (arrows) are observed on the C. scaffolds with no
  - electrospun tidemark when compared to the D. scaffolds containing an electrospun tidemark
- 17 (scale bars = A-B. 2mm, C-D. 400um).